

DOI: <http://doi.org/10.21698/simi.2023.ab17>

MAPPING OF AQUATIC ECOSYSTEMS USING GEOGRAPHIC INFORMATION SYSTEMS

Florin-Valentin Ciobotaru^{1,2}, Catalina Stoica¹

¹ National Research and Development Institute for Industrial Ecology – ECOIND, 57-73 Drumul Podu Dambovitei, district 6, 060652, Bucharest, florin.ciobotaru@ecoind.ro, Romania

²University of Bucharest, Faculty of Geography, 1 Nicolae Balcescu, 010041, Bucharest, Romania

Keywords: *aquatic ecosystems, GIS, rivers, water*

Introduction

Aquatic ecosystems include freshwater habitats such as lakes, ponds, rivers and streams, wetlands, oceans, seas which accommodate water-dependent living species including animals, plants, microbes and provide multiple benefits to humans such as *i*) provisioning services (e.g. food, water); *ii*) regulating services (that affect climate, floods, disease, wastes and water quality); *iii*) cultural services that provide recreational, aesthetic, and spiritual benefits and *iv*) supporting services such as soil formation, photosynthesis and nutrient cycling. In the last decades, continuous anthropogenic pressures were exerted on aquatic ecosystems worldwide influencing the water quality and reshaping their natural paths and areas. Early identification of pressures and understanding of the relationship between them and aquatic system could help draw management measures to achieve a good ecological status which is the main target of Water Framework Directive. Thus, in the age of technology, different application tools were developed to rapidly identify target issues on aquatic ecosystems. It is the case of the geographic information system (GIS) which is a valuable spatial system that creates, manages, analyzes and maps all types of data. The main applications that GIS could be used related to aquatic ecosystems are: *i*) *water quality assessment* (it can be used to integrate data from various sources, including remote sensing and water sampling to assess water quality in lakes, rivers and coastal areas; *ii*) *habitat mapping* (it can be used to map various aquatic habitats such as wetlands, seagrass beds, or aquatic vegetation, providing valuable information on species distribution and biodiversity); *iii*) *hydrological modeling* (it can be used to create hydrological models, to understand flow patterns, flooding risk and erosion potential in aquatic ecosystems or *iv*) *climate change impact assessment* (it can map out, model and predict the impacts of climate change on aquatic ecosystems, including changes in temperature, precipitation and sea-level rise). Moreover, a great advantage of using GIS is that it improves communication and efficiency and better management for rapidly decision making. Our study aims to apply the GIS tool for mapping two aquatic ecosystems in urban areas of Romania, namely Dambovita River and Bahlui River. The freshwater systems were digitally mapped by vectorising the level curves in order to calculate the altitudinal model and the reference surface.

Materials and methods

Studied areas. Bahlui River has a total length of 119 km, a basin area of 2,007 km² and an average annual flow of 4.88 m³/s.

Dambovita River is a water course in Romania, a tributary of the Argeş River with a total length of 237 km, a basin area of 2837 km² and an average annual flow of 19.1 m³/s.

GIS application. The design of aquatic ecosystems digital maps was done using Quantum Gis, version 3.26, Buenos Aires by QGIS Development Team. QGIS is an open-source desktop geographic information systems application that supports viewing, editing, and analyzing geospatial data. Certain digital tools such as: clicks along the hydrographic network in question for an accuracy in identifying the valleys and level curves was used for vectorization. At the same time, with the help of the coordinate reference system (CRS), the length and the shape diameter of Bahlui and Dambovita Rivers were updated. In order to frame the analyzed areas within the perimeter of Romania, the geodetic system, namely EPSG:3844 - Pulkovo 1942(58) / Stereo70 was used. In addition to all the geospatial references and digital analyses, water, soil, vegetation were represented as well.

Results and conclusions

Bahlui and Dambovita Rivers of urban areas in Romania were mapped using digital spatial system. Bahlui River runs through the Lower Jijiei Plain in the NW-SE direction, passing through the cities of Harlau and Iasi. The city of Iasi is crossed by Bahlui River over a distance of 14 km, practically dividing the city in two (Figure 1a). Dambovita River crosses the city of Bucharest for a distance of approximately 22 km, running through the city from northwest to southeast (Figure 1b). GIS allowed us to identify and map hydrological features such as watersheds and water quality monitoring sites along the two aquatic systems.




Fig.1. Spatial distribution of sampling points using GIS in **a)** Iasi (Bahlui River) and **b)** Bucharest (Dambovita River)

The digital representations of Bahlui and Dambovita Rivers showed the spatial distribution of the studied areas and leading us to identify *i)* possible pressure point sources such as hospitals and urban wastewater treatment plants (WWTPs), as well as *ii)* the sampling points upstream and downstream pressure point sources along the rivers length. The wastewater discharge into the river may influence river water quality and implicitly, the biological communities and the stability of the system as a whole.

This spatial visualisation and the data layers allowed us to assess the vulnerability of these features to environmental changes and human activities, thus guiding for possible pollution scenarios and sustainable water management solution. Although in this study only graphical representation of sampling points and the main possible pressure points were represented, the water quality results were correlated at this point only with high loading of pharmaceutical substances (PhC). Thus, we consider that the general use of pharmaceuticals compounds by urban population associated with the wastewater evacuated from hospitals alongside the lack of technologies for removal of PhC from WWTPs could be correlated with a water quality decrease.

Moreover, we assume that GIS has become an indispensable tool in hydrology, revolutionizing the field by enhancing our understanding of water processes, aiding in watershed management, and enabling effective disaster mitigation. As the world faces growing challenges related to water resources and climate change, the importance of GIS in hydrological mapping continues to grow, providing valuable insights and tools to address these pressing issues. Overall, further researches need to be carried out using GIS as a warning tool, for real-time monitoring and response during floods, droughts, as well as other hydrological events.

Acknowledgments. This work was carried out through the "Nucleu" Program within the National Research Development and Innovation Plan 2019-2022 as well as 2022-2027 with the support of Romanian Ministry of Research, Innovation and Digitalization, contract no. 20N/2019, Project code PN 19 04 02 01 and contract no.3N/2022, Project code PN 23 22 02 01.